

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use TRULANCE safely and effectively. See full prescribing information for TRULANCE.

TRULANCE® (plecanatide) tablets, for oral use  
Initial U.S. Approval: 2017

**WARNING: RISK OF SERIOUS DEHYDRATION IN PEDIATRIC PATIENTS**  
*See full prescribing information for complete boxed warning.*

- TRULANCE is contraindicated in patients less than 6 years of age; in young juvenile mice, plecanatide caused death due to dehydration. (4, 8.4)
- Avoid use of TRULANCE in patients 6 years to less than 18 years of age. (5.1, 8.4)
- The safety and effectiveness of TRULANCE have not been established in patients less than 18 years of age. (8.4)

### INDICATIONS AND USAGE

TRULANCE is a guanylate cyclase-C agonist indicated in adults for treatment of:

- chronic idiopathic constipation (CIC). (1)
- irritable bowel syndrome with constipation (IBS-C). (1)

### DOSAGE AND ADMINISTRATION

The recommended adult dosage of TRULANCE is

- CIC: 3 mg taken orally once daily. (2.1)
- IBS-C: 3 mg taken orally once daily. (2.1)

### Administration Instructions (2.2):

- Take with or without food.
- Swallow tablets whole.
- For patients who have difficulty swallowing tablets whole or those with a nasogastric or gastric feeding tube, see full prescribing information with instructions for crushing the tablet and administering with applesauce or water.

### DOSAGE FORMS AND STRENGTHS

Tablets: 3 mg (3)

### CONTRAINDICATIONS

- Patients less than 6 years of age due to the risk of serious dehydration. (4, 5.1, 8.4)
- Patients with known or suspected mechanical gastrointestinal obstruction. (4)

### WARNINGS AND PRECAUTIONS

Diarrhea: Patients may experience severe diarrhea. If severe diarrhea occurs, suspend dosing and rehydrate the patient. (5.2)

### ADVERSE REACTIONS

Most common adverse reaction ( $\geq 2\%$ ) is diarrhea. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Salix Pharmaceuticals at 1-800-321-4576 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 04/2021

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### WARNING: RISK OF SERIOUS DEHYDRATION IN PEDIATRIC PATIENTS

#### 1 INDICATIONS AND USAGE

#### 2 DOSAGE AND ADMINISTRATION

- 2.1 Recommended Dosage
- 2.2 Preparation and Administration Instructions

#### 3 DOSAGE FORMS AND STRENGTHS

#### 4 CONTRAINDICATIONS

#### 5 WARNINGS AND PRECAUTIONS

- 5.1 Risk of Serious Dehydration in Pediatric Patients
- 5.2 Diarrhea

#### 6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

#### 8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

#### 11 DESCRIPTION

#### 12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

#### 13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

#### 14 CLINICAL STUDIES

- 14.1 Chronic Idiopathic Constipation (CIC)
- 14.2 Irritable Bowel Syndrome with Constipation (IBS-C)

#### 16 HOW SUPPLIED/STORAGE AND HANDLING

#### 17 PATIENT COUNSELING INFORMATION

\*Sections or subsections omitted from the full prescribing information are not listed.

## FULL PRESCRIBING INFORMATION

**WARNING: RISK OF SERIOUS DEHYDRATION IN PEDIATRIC PATIENTS**

- TRULANCE is contraindicated in patients less than 6 years of age; in nonclinical studies in young juvenile mice, administration of a single oral dose of plecanatide caused deaths due to dehydration [see Contraindications (4), Use in Specific Populations (8.4)].
- Avoid use of TRULANCE in patients 6 years to less than 18 years of age [see Warnings and Precautions (5.1), Use in Specific Populations (8.4)].
- The safety and effectiveness of TRULANCE have not been established in patients less than 18 years of age [see Use in Specific Populations (8.4)].

### 1 INDICATIONS AND USAGE

TRULANCE is indicated in adults for the treatment of:

- chronic idiopathic constipation (CIC).
- irritable bowel syndrome with constipation (IBS-C).

### 2 DOSAGE AND ADMINISTRATION

#### 2.1 Recommended Dosage

The recommended dosage of TRULANCE for the treatment of CIC and IBS-C is 3 mg taken orally once daily.

### 2.2 Preparation and Administration Instructions

- Take TRULANCE with or without food [see Clinical Pharmacology (12.3)].
- If a dose is missed, skip the missed dose and take the next dose at the regular time. Do not take two doses at the same time.
- Swallow a tablet whole for each dose.
- For adult patients with swallowing difficulties, TRULANCE tablets can be crushed and administered orally either in applesauce or with water or administered with water via a nasogastric or gastric feeding tube. Mixing TRULANCE crushed tablets in other soft foods or in other liquids has not been tested.

#### Oral Administration in Applesauce:

1. In a clean container, crush the TRULANCE tablet to a powder and mix with 1 teaspoonful of room temperature applesauce.
2. Consume the entire tablet-applesauce mixture immediately. Do not store the mixture for later use.

#### Oral Administration in Water:

1. Place the TRULANCE tablet in a clean cup.
2. Pour approximately 30 mL of room temperature water into the cup.
3. Mix by gently swirling the tablet and water mixture for at least 10 seconds. The TRULANCE tablet will fall apart in the water.

- Swallow the entire contents of the tablet-water mixture immediately.
- If any portion of the tablet is left in the cup, add another 30 mL of water to the cup, swirl for at least 10 seconds, and swallow immediately.
- Do not store the tablet-water mixture for later use.

#### Administration with Water via a Nasogastric or Gastric Feeding Tube:

- Place the TRULANCE tablet in a clean cup with 30 mL of room temperature water.
- Mix by gently swirling the tablet and water mixture for at least 15 seconds. The TRULANCE tablet will fall apart in the water.
- Flush the nasogastric or gastric feeding tube with 30 mL of water using a catheter tip syringe.
- Draw up the mixture using the syringe and immediately administer via the nasogastric or gastric feeding tube. Do not reserve for future use.
- If any portion of the tablet is left in the cup, add another 30 mL of water to the cup, swirl for at least 15 seconds, and using the same syringe, administer via the nasogastric or gastric feeding tube.
- Using the same or a fresh syringe, flush the nasogastric or gastric feeding tube with at least 10 mL of water.

### 3 DOSAGE FORMS AND STRENGTHS

#### TRULANCE Tablets:

3 mg: white to off-white, plain, round tablet debossed with “SP” on one side and “3” for 3 mg on the other side.

### 4 CONTRAINDICATIONS

TRULANCE is contraindicated in:

- Patients less than 6 years of age due to the risk of serious dehydration [see Warnings and Precautions (5.1), Use in Specific Populations (8.4)].
- Patients with known or suspected mechanical gastrointestinal obstruction.

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Risk of Serious Dehydration in Pediatric Patients

TRULANCE is contraindicated in patients less than 6 years of age. The safety and effectiveness of TRULANCE in patients less than 18 years of age have not been established. In young juvenile mice (human age equivalent of approximately 1 month to less than 2 years), plecanatide increased fluid-secretion into the intestines as a consequence of stimulation of guanylate cyclase-C (GC-C), resulting in mortality in some mice within the first 24 hours, apparently due to dehydration. Due to increased intestinal expression of GC-C, patients less than 6 years of age may be more likely than patients 6 years of age and older to develop severe diarrhea and its potentially serious consequences.

Avoid the use of TRULANCE in patients 6 years to less than 18 years of age. Although there were no deaths in older juvenile mice, given the deaths in younger mice and the lack of clinical safety and efficacy data in pediatric patients, avoid the use of TRULANCE in patients 6 years to less than 18 years of age [see Contraindications (4), Warnings and Precautions (5.2), Use in Specific Populations (8.4)].

#### 5.2 Diarrhea

Diarrhea was the most common adverse reaction in four placebo-controlled clinical trials, two in patients with CIC and two in patients with IBS-C. Severe diarrhea was reported in 0.6% of patients in two trials in patients with CIC and in 0.6% of patients in the two trials in patients with IBS-C [see Adverse Reactions (6.1)]. If severe diarrhea occurs, suspend dosing and rehydrate the patient.

### 6 ADVERSE REACTIONS

#### 6.1 Clinical Trials Experience

Because clinical studies are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared with rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Demographic characteristics were comparable between the TRULANCE and placebo groups in all studies [see Clinical Studies (14)].

#### Chronic Idiopathic Constipation (CIC)

The safety data described below reflect data from 1,733 adult patients with CIC randomized in two double-blind, placebo-controlled clinical trials (Study 1 and Study 2) to receive placebo or 3 mg of TRULANCE once daily for 12 weeks.

#### Most Common Adverse Reactions

Table 1 provides the incidence of adverse reactions reported in at least 2% of CIC patients in the TRULANCE-treated group and at an incidence that was greater than in the placebo group.

**Table 1: Most Common Adverse Reactions<sup>a</sup> in Two Placebo-Controlled Trials of TRULANCE [Study 1 and Study 2] in Patients with CIC**

Adverse Reaction	TRULANCE, 3 mg (N = 863) %	Placebo (N = 870) %
Diarrhea <sup>b</sup>	5	1

<sup>a</sup> Reported in at least 2% of TRULANCE-treated patients with CIC and at an incidence greater than placebo.

<sup>b</sup> Verbatim reports of diarrhea were recorded as adverse reactions; reports of loose stools and increase in stool frequency were recorded as adverse reactions if they were also reported to be bothersome to the patient.

#### Diarrhea

The majority of reported cases of diarrhea occurred within 4 weeks of treatment initiation. Severe diarrhea was reported in 0.6% of TRULANCE-treated patients compared to 0.3% of placebo-treated patients. Severe diarrhea was reported to occur within the first 3 days of treatment [see Warnings and Precautions (5.2)].

#### Adverse Reactions Leading to Discontinuation

Discontinuations due to adverse reactions occurred in 4% of TRULANCE-treated patients and 2% of placebo-treated patients. The most common adverse reaction leading to discontinuation was diarrhea: 2% of TRULANCE-treated patients and 0.5% of placebo-treated patients withdrew due to diarrhea.

#### Less Common Adverse Reactions

Adverse reactions reported in less than 2% of TRULANCE-treated patients and at an incidence greater than placebo were: sinusitis, upper respiratory tract infection, abdominal distension, flatulence, abdominal tenderness, and increased liver biochemical tests (2 patients with alanine aminotransferase (ALT) greater than 5 to 15 times the upper limit of normal and 3 patients with aspartate aminotransferase (AST) greater than 5 times the upper limit of normal).

#### Irritable Bowel Syndrome with Constipation (IBS-C)

The safety data described below reflect data from 1,449 adults patients with IBS-C randomized in two double-blind, placebo-controlled clinical trials (Study 3 and Study 4) to receive placebo or 3 mg TRULANCE once daily for 12 weeks.

#### Most Common Adverse Reactions

Table 2 provides the incidence of adverse reactions reported in at least 2% of IBS-C patients treated with TRULANCE and at an incidence that was greater than in the placebo group.

**Table 2: Most Common Adverse Reactions<sup>a</sup> in Two Placebo-Controlled Trials of TRULANCE [Study 3 and Study 4] in Patients with IBS-C**

Adverse Reaction	TRULANCE, 3 mg (N = 723) %	Placebo (N = 726) %
Diarrhea <sup>b</sup>	4.3	1

<sup>a</sup> Reported in at least 2% of TRULANCE-treated patients with IBS-C and at an incidence greater than placebo.

<sup>b</sup> Verbatim reports of diarrhea were recorded as adverse reactions; reports of loose stools and increase in stool frequency were recorded as adverse reactions if they were also reported to be bothersome to the patient.

#### Diarrhea

The majority of reported cases of diarrhea occurred within 4 weeks of treatment initiation. Severe diarrhea was reported in 1% of TRULANCE-treated patients compared to 0.1% of placebo-treated patients [see Warnings and Precautions (5.2)]. Severe diarrhea was reported to occur within the first day of treatment.

#### Adverse Reactions Leading to Discontinuation

Discontinuations due to adverse reactions occurred in 2.5% of TRULANCE-treated patients and 0.4% of placebo-treated patients. The most common adverse reaction leading to discontinuation was diarrhea: 1.2% of TRULANCE-treated patients and 0% of placebo-treated patients withdrew due to diarrhea.

#### Less Common Adverse Reactions

Adverse reactions reported in 1% or more but less than 2% of TRULANCE-treated patients and at an incidence greater than placebo were: nausea, nasopharyngitis, upper respiratory tract infection, urinary tract infection, and dizziness. Two patients reported increased liver biochemical tests (alanine aminotransferase (ALT) greater than 5 to 15 times the upper limit of normal).

### 6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of TRULANCE. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to TRULANCE exposure.



## Elimination

### Metabolism

Plecanatide was metabolized in the GI tract to an active metabolite by loss of the terminal leucine moiety. Both plecanatide and the metabolite were proteolytically degraded within the intestinal lumen to smaller peptides and naturally occurring amino acids.

### Excretion

No excretion studies have been conducted in humans. Plecanatide and its active metabolite were not measurable in plasma following administration of the recommended clinical doses.

### Drug Interaction Studies

Neither plecanatide nor its active metabolite inhibited the cytochrome P450 (CYP) enzymes 2C9 and 3A4, and they did not induce CYP3A4 *in vitro*.

Plecanatide and its active metabolite were neither substrates nor inhibitors of the transporters P-glycoprotein (P-gp) or breast cancer resistance protein (BCRP) *in vitro*.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

#### Carcinogenesis

The carcinogenic potential of plecanatide was assessed in 2-year carcinogenicity studies in mice and rats. Plecanatide was not tumorigenic in mice at oral doses up to 90 mg/kg/day or in rats at oral doses up to 100 mg/kg/day. Limited systemic exposure to plecanatide was achieved at the tested dose levels in animals, whereas no detectable exposure occurred in humans. Therefore, animal and human doses should not be compared directly for evaluating relative exposure.

#### Mutagenesis

Plecanatide was not genotoxic in the *in vitro* bacterial reverse mutation (Ames) assay, *in vitro* mouse lymphoma mutation assay, or the *in vivo* mouse bone marrow micronucleus assay.

#### Impairment of Fertility

Plecanatide had no effect on fertility or reproductive function in male or female mice at oral doses of up to 600 mg/kg/day.

## 14 CLINICAL STUDIES

### 14.1 Chronic Idiopathic Constipation (CIC)

The efficacy of TRULANCE for the management of symptoms of CIC was established in two 12-week, double-blind, placebo-controlled, randomized, multicenter clinical studies in adult patients (Study 1 and Study 2). In the Intention-to-Treat (ITT) population, a total of 905 patients (Study 1) and 870 patients (Study 2) were randomized 1:1 to either placebo or TRULANCE 3 mg, once daily. In clinical studies, study medication was administered without respect to food intake. Demographics for these studies included an overall mean age of 45 years (range 18 to 80 years), 80% female, 72% white, and 24% black.

To be eligible for the studies, patients were required to meet modified Rome III criteria for at least 3 months prior to the screening visit, with symptom onset for at least 6 months prior to diagnosis. Rome III criteria were modified to require that patients report less than 3 defecations per week, rarely have a loose stool without the use of laxatives, not use manual maneuvers to facilitate defecations, and not meet criteria for IBS-C. In addition, patients were required to report at least two of the following symptoms:

- Straining during at least 25% of defecations
- Lumpy or hard stool in at least 25% of defecations
- Sensation of incomplete evacuations for at least 25% of defecations
- Sensation of anorectal obstruction/blockage for at least 25% of defecations

Patients who met these criteria were also required to demonstrate the following during the last 2 weeks of the screening period:

- Less than 3 complete spontaneous bowel movements (CSBMs) (a CSBM is an SBM that is associated with a sense of complete evacuation) in each of the two weeks
- Bristol Stool Form Scale (BSFS) of 6 or 7 in less than 25% of spontaneous bowel movements (SBMs) (an SBM is a bowel movement occurring in the absence of laxative use)
- One out of the following three:
  - BSFS of 1 or 2 in at least 25% of defecations
  - A straining value recorded on at least 25% of days when a BM was reported
  - At least 25% of BMs result in a sense of incomplete evacuation

The efficacy of TRULANCE was assessed using a responder analysis and change-from-baseline in CSBM and SBM endpoints. Efficacy was assessed using information provided by patients on a daily basis in an electronic diary.

A responder was defined as a patient who had at least 3 CSBMs in a given week and an increase of at least 1 CSBM from baseline in the same week for at least 9 weeks out of the 12-week treatment period and at least 3 of the last 4 weeks of the study. The responder rates are shown in Table 3.

**Table 3: Efficacy Responder Rates in the Two Placebo-controlled Studies of CIC: at least 9 of 12 weeks and at least 3 of the last 4 weeks (ITT Population)**

Study 1			
	TRULANCE 3 mg N = 453	Placebo N = 452	Treatment Difference <sup>a</sup> [95% CI <sup>b</sup> ]
Responder <sup>c</sup>	21%	10%	11% [6.1%, 15.4%]
Study 2			
	TRULANCE 3 mg N = 430	Placebo N = 440	Treatment Difference <sup>a</sup> [95% CI <sup>b</sup> ]
Responder <sup>c</sup>	21%	13%	8% [2.6%, 12.4%]

<sup>a</sup> p-value <0.005

<sup>b</sup> CI = confidence interval

<sup>c</sup> Primary endpoint defined as a patient who had at least 3 CSBMs in a given week and an increase of at least 1 CSBM from baseline in the same week for at least 9 weeks out of the 12-week treatment period and at least 3 of the last 4 weeks of the study.

In both studies, improvements in the frequency of CSBMs/week were seen as early as week 1 with improvement maintained through week 12. The difference between the TRULANCE group and the placebo group in the mean change of CSBMs/week frequency from baseline to week 12 was approximately 1.1 CSBMs/week.

Over the 12-week treatment period, improvements were observed in stool frequency (number of CSBMs/week and SBMs/week) and/or stool consistency (as measured by the BSFS), and/or in the amount of straining with bowel movements (amount of time pushing or physical effort to pass stool) in the TRULANCE group as compared to placebo.

Following completion of the study drug treatment period, patients continued to record data in the daily diary for a 2-week Post-Treatment Period. During this time, TRULANCE-treated patients generally returned to baseline for these study endpoints.

In Studies 1 and 2, a third randomized treatment arm of TRULANCE 6 mg once daily did not demonstrate additional treatment benefit and had a greater incidence of adverse reactions than TRULANCE 3 mg once daily. Therefore, TRULANCE 6 mg once daily is not recommended [see *Dosage and Administration* (2.1)].

### 14.2 Irritable Bowel Syndrome with Constipation (IBS-C)

The efficacy of TRULANCE for the management of symptoms of IBS-C was established in two 12-week, double-blind, placebo-controlled, randomized, multicenter clinical studies in adult patients (Study 3 and Study 4). In the Intention-to-Treat (ITT) population, a total of 699 patients (Study 3) and 754 patients (Study 4) received treatment with placebo or TRULANCE 3 mg once daily. In clinical studies, study medication was administered without respect to food intake. Demographics for these studies included an overall mean age of 44 years (range 18 to 83 years), 74% female, 73% white, and 22% black.

To be eligible, patients were required to meet the Rome III criteria for IBS for at least 3 months prior to the screening visit, with symptom onset for at least 6 months prior to diagnosis. Diagnosis required recurrent abdominal pain or discomfort at least 3 days/month in the last 3 months associated with 2 or more of 1) improvement with defecation, 2) onset associated with a change in frequency of stool, and 3) onset associated with a change in form (appearance) of stool. Patients also met the IBS-C differentiation criteria for constipation, characterized by a stool pattern such that at least 25% of defecations are hard or lumpy stools and no more than 25% of defecations are loose or watery stool.

Patients who met these criteria were excluded if they demonstrated the following during the last 2 weeks of the screening period:

- Worst abdominal pain intensity (WAPI) score of 0 on an 11-point scale for more than 2 days during each week
- An average WAPI of less than 3 for either week
- More than 3 complete spontaneous bowel movements (CSBMs) or more than six spontaneous bowel movements (SBMs) per week in either week
- Bristol Stool Form Scale (BSFS) of 7 for any SBM recorded
- More than 1 day in either week with a BSFS of 6 for any SBM recorded
- No use of rescue laxative (bisacodyl) within 72 hours before randomization

The efficacy of TRULANCE was assessed using a responder analysis based on abdominal pain intensity and a stool frequency responder (CSBM) endpoint. Efficacy was assessed using information provided by patients on a daily basis through an electronic phone diary system.

A responder was defined as a patient who met both the abdominal pain intensity and stool frequency responder criteria in the same week for at least 6 of the 12 treatment weeks. The abdominal pain intensity and stool frequency responder criteria assessed each week were defined as:

- Abdominal pain intensity responder: a patient who experienced a decrease in the weekly average of worst abdominal pain in the past 24 hours score (measured daily) of at least 30% compared with baseline weekly average.
- Stool frequency responder: a patient who experienced an increase of at least 1 CSBM per week from baseline.

The responder rates are shown in Table 4.

**Table 4: Efficacy Responder Rates in the Two Placebo-controlled Studies of IBS-C: Overall Responder for at least 6 of the 12 Treatment Weeks (ITT Population)**

Study 3			
	Placebo N = 350	TRULANCE 3 mg N = 349	Treatment Difference [95% CI <sup>a</sup> ]
<b>Responder<sup>b</sup></b>	18%	30%	12% [6%, 18%]
<u>Components of Responder Endpoint</u>			
Abdominal Pain Responder <sup>c</sup>	32%	41%	
CSBM Responder <sup>d</sup>	35%	48%	
Study 4			
	Placebo N = 379	TRULANCE 3 mg N = 375	Treatment Difference [95% CI <sup>a</sup> ]
<b>Responder<sup>b</sup></b>	14%	21%	7% [2%, 13%]
<u>Components of Responder Endpoint</u>			
Abdominal Pain Responder <sup>c</sup>	23%	33%	
CSBM Responder <sup>d</sup>	28%	34%	

<sup>a</sup>: CI = confidence interval

<sup>b</sup>: A responder for these trials was defined as a patient who met both the abdominal pain and CSBM weekly responder criteria for at least 6 of the 12 weeks.

<sup>c</sup>: An abdominal pain responder was defined as a patient who met the criteria of at least 30% reduction from baseline in weekly average of the worst daily abdominal pain, for at least 6 of the 12 weeks.

<sup>d</sup>: A CSBM responder was defined as a patient who achieved an increase in at least 1 CSBM per week, from baseline, for at least 6 of 12 weeks.

In both studies, the proportion of responders who were also weekly responders for at least 2 of the 4 treatment weeks in month 3, the last month of treatment was greater in the TRULANCE groups compared to placebo.

Over the 12-week treatment period, improvements were observed in both stool consistency (as measured by the BSFS) and in the amount of straining with bowel movements (amount of time pushing or physical effort to pass stool) in the 3 mg TRULANCE group as compared to placebo.

Following completion of the study drug treatment period, patients continued to record data in the daily diary for a 2-week Post-Treatment Period. During this time, TRULANCE-treated patients generally returned to baseline for these study endpoints.

In Studies 3 and 4, a third randomized treatment arm of TRULANCE 6 mg once daily did not demonstrate additional treatment benefit over the 3 mg dose. Therefore, TRULANCE 6 mg once daily is not recommended [see *Dosage and Administration* (2.1)].

## 16 HOW SUPPLIED/STORAGE AND HANDLING

TRULANCE tablets are packaged in an aluminum foil unit dose blister pack of 30 in a child-resistant pack or in a white, opaque, high-density polyethylene round bottle with a screw-top polypropylene child-resistant cap and heat-activated induction seal. Each bottle container-closure system also contains a desiccant and a polyester coil.

TRULANCE 3 mg tablets are white to off-white, plain and round, debossed with “SP” on one side and “3” for 3 mg on the other side and supplied as:

NDC Number	Size
65649-003-30	Bottle of 30
70194-003-30	Aluminum foil unit dose blister pack of 30 in a child-resistant pack

Store at room temperature, 20 to 25°C (68 to 77°F); excursions permitted to 15 to 30°C (59 to 86°F) [see USP Controlled Room Temperature].

Keep TRULANCE in a dry place. Protect from moisture. For bottles, keep TRULANCE in the original bottle. Do not remove desiccant from the bottle. Do not subdivide or repack.

## 17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

Advise Patients:

### Diarrhea

To stop TRULANCE and contact their healthcare provider if they experience severe diarrhea [see *Warnings and Precautions* (5.2)].

### Accidental Ingestion

Accidental ingestion of TRULANCE in children, especially in children less than 6 years of age, may result in severe diarrhea and dehydration. Instruct patients to take steps to store TRULANCE securely and out of reach of children and to dispose of unused TRULANCE [see *Contraindications* (4), *Warnings and Precautions* (5.2)].

### Administration and Handling Instructions

- To take TRULANCE once daily with or without food [see *Dosage and Administration* (2.2)].
- If a dose is missed, skip the missed dose and take the next dose at the regular time. Do not take two doses at the same time.
- To swallow TRULANCE tablets whole.
- If adult patients have swallowing difficulties, TRULANCE tablets can be crushed and administered orally in either applesauce or with water, or administered with water via a nasogastric or gastric feeding tube, as described in the Medication Guide.
- To keep TRULANCE in a dry place. Protect from moisture. For bottles, keep TRULANCE in the original bottle. Do not remove desiccant from the bottle. Do not subdivide or repack. Remove and discard polyester coil after opening. Keep bottles closed tightly [see *How Supplied/Storage and Handling* (16)].

### Distributed by:

Salix Pharmaceuticals, a division of  
Bausch Health US, LLC  
Bridgewater, NJ 08807 USA

U.S. Patent Numbers: 7,041,786; 7,799,897; 8,637,451; 9,610,321; 9,616,097; 9,919,024; 9,925,231 and 10,011,637

TRULANCE is a trademark of Salix Pharmaceuticals, Inc. or its affiliates.

© 2021 Salix Pharmaceuticals, Inc. or its affiliates

9705703

**Medication Guide**  
**TRULANCE® (TROO lans)**  
**(plecanatide) tablets, for oral use**

**What is the most important information I should know about TRULANCE?**

- **Do not give TRULANCE to children who are less than 6 years of age. It may harm them.**
- **You should not give TRULANCE to children 6 years to less than 18 years of age. It may harm them.**

See “**What are the possible side effects of TRULANCE?**” for more information about side effects.

**What is TRULANCE?**

TRULANCE is a prescription medicine used in adults to treat:

- a type of constipation called chronic idiopathic constipation (CIC). Idiopathic means the cause of the constipation is unknown.
- irritable bowel syndrome with constipation (IBS-C).

**It is not known if TRULANCE is safe and effective in children less than 18 years of age.**

**Who should not take TRULANCE?**

- **Do not give TRULANCE to children who are less than 6 years of age.**
- Do not take TRULANCE if a doctor has told you that you have a bowel blockage (intestinal obstruction).

**Before taking TRULANCE, tell your doctor about all of your medical conditions, including if you:**

- are pregnant or plan to become pregnant. It is not known if TRULANCE will harm your unborn baby.
- are breastfeeding or plan to breastfeed. Talk with your doctor about the best way to feed your baby if you take TRULANCE.

**Tell your doctor about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**How should I take TRULANCE?**

- Take TRULANCE exactly as your doctor tells you to take it.
- Take TRULANCE by mouth, 1 time each day with or without food.
- If you miss a dose, skip the missed dose. Take the next dose at your regular time. Do not take 2 doses at the same time.
- TRULANCE tablets should be swallowed whole.

Adults who cannot swallow TRULANCE tablets whole may crush the TRULANCE tablet and mix with applesauce or dissolve TRULANCE in water before swallowing. TRULANCE tablets may also be taken with water by adults through a nasogastric or gastric feeding tube.

It is not known if TRULANCE is safe and effective when crushed and mixed with other foods or dissolved in other liquids.

**Taking TRULANCE in applesauce:**

- Crush the TRULANCE tablet in a clean container until it is a powder and mix with 1 teaspoon of room temperature applesauce.
- Swallow all of the TRULANCE and applesauce mixture right away. Do not keep the TRULANCE and applesauce mixture for future use.

**Taking TRULANCE in water:**

- Place the TRULANCE tablet in a clean cup and pour 1 ounce (30 mL) of room temperature water into the cup.
- Gently swirl the TRULANCE tablet and water for at least 10 seconds. The TRULANCE tablet will fall apart in the water.
- Swallow all of the TRULANCE tablet and water mixture right away. Do not keep the mixture for future use.
- If you see any part of the tablet left in the cup, add another 1 ounce (30 mL) of water to the cup, swirl for at least 10 seconds, and swallow right away.

**Taking TRULANCE through a nasogastric or gastric feeding tube:**

Gather the supplies you will need to take your TRULANCE dose. Your doctor should tell you what size catheter tip syringe you will need for your dose. Ask your doctor if you have any questions about how to give TRULANCE the right way.

- Place the TRULANCE tablet in a clean cup with 1 ounce (30 mL) of room temperature water.
- Gently swirl the TRULANCE tablet and water for at least 15 seconds. The TRULANCE tablet will fall apart in the water.
- Flush the nasogastric or gastric feeding tube with 1 ounce (30 mL) of water.
- Draw up the TRULANCE tablet and water mixture into a catheter tip syringe and give right away through the nasogastric or gastric feeding tube. Do not keep the mixture for future use.
- If you see any part of the tablet left in the cup, add another 1 ounce (30 mL) of water to the cup, swirl for at least 15 seconds and use the same catheter tip syringe to give the mixture through the nasogastric or gastric feeding tube.
- Using the same or another catheter tip syringe, flush the nasogastric or gastric feeding tube with at least 10 mL of water.

**What are the possible side effects of TRULANCE?**

**TRULANCE can cause serious side effects, including:**

- See “What is the most important information I should know about TRULANCE?”
- **Diarrhea is the most common side effect of TRULANCE, and it can sometimes be severe.**
  - Diarrhea often begins within the first 4 weeks of TRULANCE treatment.

**Stop taking TRULANCE and call your doctor if you develop severe diarrhea.**

These are not all the possible side effects of TRULANCE.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store TRULANCE?**

- Store TRULANCE at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep TRULANCE in a secure place and in the bottle or blister pack that it comes in.
- The TRULANCE bottle contains a desiccant packet to help keep your medicine dry (protect it from moisture). Do not remove the desiccant packet from the bottle.
- The TRULANCE bottle contains a polyester coil to help protect the tablets during shipping. Remove the polyester coil from the bottle and throw it away after opening the bottle.
- Keep the container of TRULANCE tightly closed and in a dry place.
- Safely throw away TRULANCE that is out of date or no longer needed.

**Keep TRULANCE and all medicines out of the reach of children.**

**General information about the safe and effective use of TRULANCE.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use TRULANCE for a condition for which it was not prescribed. Do not give TRULANCE to other people, even if they have the same symptoms that you have. It may harm them.

You can ask your doctor or pharmacist for information about TRULANCE that is written for health professionals.

**What are the ingredients in TRULANCE?**

**Active ingredient:** plecanatide

**Inactive ingredients:** magnesium stearate and microcrystalline cellulose

**Distributed by:**

Salix Pharmaceuticals, a division of  
Bausch Health US, LLC  
Bridgewater, NJ 08807 USA

U.S. Patent Numbers: 7,041,786; 7,799,897; 8,637,451; 9,610,321; 9,616,097; 9,919,024; 9,925,231 and 10,011,637

TRULANCE is a trademark of Salix Pharmaceuticals, Inc. or its affiliates.

© 2021 Salix Pharmaceuticals, Inc. or its affiliates

This Medication Guide is approved by the U.S. Food and Drug Administration.

02/2021

9705703  
REF-TRU-0381